

Recommended reading material for the understanding of iron ore deposits

in blue: CET and UFMG publications

in bold: recommended reading by the Iron Research Group, CET. In most cases, these recommended articles are reviews of the topic and therefore represent a good starting point for reading.

Banded Iron Formations - genesis and characteristics through space and time

Barley, M. E., Pickard, A. L., and Sylvester, P. J., 1997, Emplacement of a large igneous province as a possible cause of banded iron formation 2.45 billion years ago: *Nature*, v. 385, p. 55-58.

Bekker, A., Slack, J. F., Planavsky, N., Krapez, B., Hofmann, A., Konhauser, K. O., and Rouxel, O. J., 2010, Iron Formation: The Sedimentary Product of a Complex Interplay among Mantle, Tectonic, Oceanic, and Biospheric Processes: *Economic Geology*, v. 105, p. 467-508.

Beukes, N. J., and Gutzmer, J., 2008, Origin and Paleoenvironmental Significance of Major Iron Formations at the Archean-Paleoproterozoic Boundary, in Hagemann, S. G., Rosière, C. A., Gutzmer, J., and Beukes, N. J., eds., *Banded Iron Formation-related High-grade Iron Ore. Reviews in Economic Geology*, v. 15, p. 5-47.

Czaja, A. D., Johnson, C. M., Beard, B. L., and Van Kranendonk, M. J., 2010, Iron isotope evidence for an abiological origin of a BIF from the Yilgarn Craton: 5th International Archean Symposium: Abstracts, Perth, Western Australia, 2010, p. 349-351.

Gole, M. J., 1980, Mineralogy and petrology of very-low-metamorphic grade Archaean banded iron-formations, Weld Range, Western Australia: *American Mineralogist*, v. 65, p. 1-2.

---, 1981, Archean Banded Iron-Formations, Yilgarn Block, Western Australia: *Economic Geology*, v. 76, p. 1954-1974.

Hagemann, S., Rosiere, C. A., Gutzmer, J., and Beukes, N. J., 2008, Glossary of terms Banded Iron Formation-related high-grade iron ore: *Banded Iron Formation-Related High-Grade Iron Ore*, v. 15, p. 411-414.

Hallimond, A. F., 1925, Iron ores: bedded ores of England and Wales. Petrography and chemistry: Memoirs of the Geological Survey of Great Britain (Special reports on the mineral resources of Great Britain), v. 29, p. 111.

Huberty, J. M., Kita, N. T., Kozdon, R., Heck, P. R., Fournelle, J. H., Spicuzza, M. J., Xu, H. F., and Valley, J. W., 2010, Crystal orientation effects in delta O-18 for magnetite and hematite by SIMS: *Chemical Geology*, v. 276, p. 269-283.

Hyslop, E. V., Valley, J. W., Johnson, C. M., and Beard, B. L., 2008, The effects of metamorphism on O and Fe isotope compositions in the Biwabik Iron Formation, northern Minnesota: *Contributions to Mineralogy and Petrology*, v. 155, p. 313-328.

Isley, A. E., and Abbott, D. H., 1999, Plume-related mafic volcanism and the deposition of banded iron formation: *J. Geophys. Res.*, v. 104, p. 15461-15477.

Klein, C., and Beukes, N. J., 1993, Sedimentology and Geochemistry of the Glaciogenic Late Proterozoic Rapitan Iron-Formation in Canada: *Economic Geology and the Bulletin of the Society of Economic Geologists*, v. 88, p. 542-565.

Konhauser, K. O., Hamade, T., Raiswell, R., Morris, R. C., Ferris, F. G., Southam, G., and Canfield, D. E., 2002, Could bacteria have formed the Precambrian banded iron formations?: *Geology*, v. 30, p. 1079-1082.

Konhauser, K. O., Newman, D. K., and Kappler, A., 2005, The potential significance of microbial Fe(III) reduction during deposition of Precambrian banded iron formations: *Geobiology*, v. 3, p. 167-177.

Krapez, B., Barley, M. E., and Pickard, A. L., 2003, Hydrothermal and resedimented origins of the precursor sediments to banded iron formation: sedimentological evidence from the Early Palaeoproterozoic Brockman Supersequence of Western Australia: *Sedimentology*, v. 50, p. 979-1011.

[Lascelles, D. F., 2007, Black smokers and density currents: A uniformitarian model for the genesis of banded iron-formations: *Ore Geology Reviews*, v. 32, p. 381-411.](#)

Pickard, A. L., Barley, M. E., and Krapež, B., 2006, Deep-marine depositional setting of banded iron formation: sedimentological evidence from interbedded clastic sedimentary rocks in the early Palaeoproterozoic Dales Gorge Member of Western Australia: *Sedimentary Geology*, v. 170, p. 37-62.

Planavsky, N., Bekker, A., Rouxel, O. J., Kamber, B., Hofmann, A., Knudsen, A., and Lyons, T. W., 2010, Rare Earth Element and yttrium compositions of Archean and Paleoproterozoic Fe formations revisited: New perspectives on the significance and mechanisms of deposition: *Geochimica et Cosmochimica Acta*, v. 74, p. 6387-6405.

Rasmussen, B., Fletcher, I. R., Muhling, J. R., Thorne, W. S., and Broadbent, G. C., 2007, Prolonged history of episodic fluid flow in giant hematite ore bodies: Evidence from in situ U-Pb geochronology of hydrothermal xenotime: *Earth and Planetary Science Letters*, v. 258, p. 249-259.

Rasmussen, B., Fletcher, I. R., Muhling, J. R., and Wilde, S. A., 2010, In situ U-Th-Pb geochronology of monazite and xenotime from the Jack Hills belt: Implications for the age of deposition and metamorphism of Hadean zircons: *Precambrian Research*, v. 180, p. 26-46.

Thurston, P. C., Kamber, B. S., and Whitehouse, M., in press, Archean cherts in banded iron formation: Insight into Neoproterozoic ocean chemistry and depositional processes: *Precambrian Research*.

Genetic models for iron ore

Barley, M. E., Pickard, A. L., Hagemann, S. G., and Folkert, S. L., 1999, Hydrothermal origin for the 2 billion year old Mount Tom Price giant iron ore deposit, Hamersley Province, Western Australia: *Mineralium Deposita*, v. 34, p. 784-789.

Clout, J. M. F., 2003, Upgrading processes in BIF-derived iron ore deposits: implications for ore genesis and downstream mineral processing: *Transactions of the Institution of Mining and Metallurgy Section B-Applied Earth Science*, v. 112, p. B89-B95.

Dalstra, H., and Guedes, S. T., 2004, Giant hydrothermal hematite deposits with Mg-Fe metasomatism: A comparison of the Carajas, Hamersley, and other iron ores: *Economic Geology*, v. 99, p. 1793-1800.

Dalstra, H., Harding, T., Riggs, T., and Taylor, D., 2003, Banded iron formation hosted high-grade hematite deposits, a coherent group?: *Transactions of the Institution of Mining and Metallurgy Section B-Applied Earth Science*, v. 112, p. B68-B72.

Dalstra, H. J., and Rosière, C. A., 2008, Structural Controls on High-Grade Iron Ores Hosted by Banded Iron Formation: A Global Perspective, in Hagemann, S., Rosière, C. A., Gutzmer, J., and Beukes, N. J., eds., *Banded Iron Formation-related High-grade Iron Ore. Reviews in Economic Geology*, v. 15, *Society of Economic Geologists*, p. 73-106.

Gruner, J. W., 1937, Hydrothermal leaching of iron ores of the Lake Superior type - a modified theory: *Economic Geology*, v. 32, p. 121-130.

Gutzmer, J., Chisonga, C. B., Beukes, N. J., and Mukhopadhyay, J., 2008, The Geochemistry of Iron Formation-Hosted High-Grade Hematite-Martite Iron Ores, in Hagemann, S. G., Rosière, C. A., Gutzmer, J., and Beukes, N. J., eds., *Banded Iron Formation-related High-grade Iron Ore. Reviews in Economic Geology*, v. 15, p. 157-183.

Gutzmer, J., Mukhopadhyay, J., Beukes, N. J., Pack, A., Hayashi, K., and Sharp, Z. D., 2006, Oxygen isotope composition of hematite and genesis of high-grade BIF-hosted iron ores: *Geological Society of America Memoir*, v. 198, p. 257-268.

Hagemann, S. G., Barley, M. E., Folkert, S. L., Yardley, B. W. D., and Banks, D. A., 1999, A hydrothermal origin for the giant BIF-hosted Tom Price iron ore deposit: *Mineral Deposits: Processes to Processing*, Vols 1 and 2, p. 41-44.

Lobato, L. M., Silva, R. C. F. E., Hagemann, S., Thorne, W., and Zucchetri, M., 2008, Hypogene Alteration Associated with High-Grade Banded Iron Formation-Related Iron Ore: *Banded Iron Formation-Related High-Grade Iron Ore*, v. 15, p. 107-128.

McLellan, J. G., Oliver, N. H. S., Ord, A., Zhang, Y., and Schaub, P. M., 2003, A numerical modelling approach to fluid flow in extensional environments: implications for genesis of large microplaty hematite ores: *Journal of Geochemical Exploration*, v. 78-79, p. 675-679.

Morris, R. C., 1985, Genesis of iron ore in banded iron-formation by supergene and supergene-metamorphic processes - a conceptual model, in Wolf, K. H., ed., *Handbook of strata-bound and stratiform ore deposits*: Amsterdam, Elsevier, p. 73-235.

Morris, R. C., and Kneeshaw, M., 2011, Genesis modelling for the Hamersley BIF-hosted iron ores of Western Australia: a critical review: *Australian Journal of Earth Sciences*, v. 58, p. 417-451.

Morris, R. C., Thornber, M. R., and Ewers, W. E., 1980, Deep-seated iron ores from banded iron-formation: *Nature*, v. 288, p. 250-252.

Powell, C. M., Oliver, N. H. S., Li, Z. X., Martin, D. M., and Ronaszeki, J., 1999, Synorogenic hydrothermal origin for giant Hamersley iron oxide ore bodies: *Geology*, v. 27, p. 175-178.

Ramanaidou, E. R., 2009, Genesis of lateritic iron ore from banded iron-formation in the Capanema mine (Minas Gerais, Brazil): *Australian Journal of Earth Sciences*, v. 56, p. 605-620.

Ramanaidou, E. R., and Morris, R. C., 2009, Comparison of Supergene Mimetic and Supergene Lateritic Iron Ore Deposits, in AusIMM, ed., *Proceedings Iron Ore Conference 2009*, 27-29 July 2009, Perth, Western Australia, p. 243-246.

Taylor, D., Dalstra, H. J., Harding, A. E., Broadbent, G. C., and Barley, M. E., 2001, Genesis of high-grade hematite orebodies of the Hamersley Province, Western Australia: *Economic Geology*, v. 96, p. 837-873.

Iron ore deposits throughout the World

Hamersley Province, Western Australia

Clout, J. M. F., 2003, Upgrading processes in BIF-derived iron ore deposits: implications for ore genesis and downstream mineral processing: Transactions of the Institution of Mining and Metallurgy Section B-Applied Earth Science, v. 112, p. B89-B95.

Clout, J. M. F., 2005, Iron formation-hosted iron ores in the Hamersley Province of Western Australia, Iron Ore Conference 2005, 9–19.

Dalstra, H., and Guedes, S. T., 2004, Giant hydrothermal hematite deposits with Mg-Fe metasomatism: A comparison of the Carajas, Hamersley, and other iron ores: Economic Geology, v. 99, p. 1793-1800.

Greentree, M. R., and Lord, D., 2007, Iron mineralization in the Yilgarn craton and future potential, in Bierlein, F. P., and Knox-Robinson, C. M., eds., Proceedings of Geoconferences (WA) Inc Kalgoorlie '07 Conference: Kalgoorlie, p. 70-73.

Haest, M., Cudahy, T., Laukamp, C., and Gregory, S., 2012, Quantitative mineralogy from infrared spectroscopic data: (I) Validation of mineral abundance and composition scripts at the Rocklea channel iron deposit in Western Australia: Economic Geology, p. 209-228.

---, 2012, Quantitative mineralogy from infrared spectroscopic data: (II) Three-dimensional mineralogical characterization of the Rocklea channel iron deposit, Western Australia: Economic Geology, v. 107, p. 229-249.

[Hagemann, S. G., Barley, M. E., Folkert, S. L., Yardley, B. W. D., and Banks, D. A., 1999, A hydrothermal origin for the giant BIF-hosted Tom Price iron ore deposit: Mineral Deposits: Processes to Processing, Vols 1 and 2, p. 41-44.](#)

Li, Z. X., Powell, C. M., and Bowman, R., 1993, Timing and genesis of Hamersley iron-ore deposits: Exploration Geophysics, v. 24, p. 631-636.

Muller, S. G., Krapez, B., Barley, M. E., and Fletcher, I. R., 2005, Giant iron-ore deposits of the Hamersley province related to the breakup of Paleoproterozoic Australia: New insights from in situ SHRIMP dating of baddeleyite from mafic intrusions: Geology, v. 33, p. 577-580.

Powell, C. M., Oliver, N. H. S., Li, Z. X., Martin, D. M., and Ronaszeki, J., 1999, Synorogenic hydrothermal origin for giant Hamersley iron oxide ore bodies: Geology, v. 27, p. 175-178.

Taylor, D., Dalstra, H. J., Harding, A. E., Broadbent, G. C., and Barley, M. E., 2001, Genesis of high-grade hematite orebodies of the Hamersley Province, Western Australia: Economic Geology, v. 96, p. 837-873.

Thorne, W., Hagemann, S., Vennemann, T., and Oliver, N., 2009, Oxygen Isotope Compositions of Iron Oxides from High-Grade BIF-Hosted Iron Ore Deposits of the Central Hamersley Province Western Australia: Constraints on the Evolution of Hydrothermal Fluids: *Economic Geology*, v. 104, p. 1019-1035.

Thorne, W., Hagemann, S., Webb, A., and Clout, J., 2008, Banded Iron Formation-Related Iron Ore Deposits of the Hamersley Province, Western Australia: Banded Iron Formation-Related High-Grade Iron Ore, v. 15, p. 197-221.

Thorne, W. S., Hagemann, S. G., and Barley, M., 2004, Petrographic and geochemical evidence for hydrothermal evolution of the North Deposit, Mt Tom Price, Western Australia: *Mineralium Deposita*, v. 39, p. 766-783.

Webb, A. D., Dickens, G. R., and Oliver, N. H. S., 2004, Carbonate alteration of the Upper Mount McRae Shale beneath the martite-microplaty hematite ore deposit at Mount Whaleback, Western Australia: *Mineralium Deposita*, v. 39, p. 632-645.

Yilgarn Craton, Western Australia

Angerer, T., Duuring, P., Lascelle, D. F., and Hagemann, S. G., 2010, BIF-related iron ore in the Yilgarn Craton, Western Australia: Geological setting and ore forming processes: 5th International Archean Symposium: Abstracts, Perth, Western Australia, 2010, p. 277-280.

Angerer, T., and Hagemann, S. G., 2010, The BIF-hosted high-grade iron ore deposits in the Archean Koolyanobbing greenstone belt, Western Australia: Structural control on synorogenic- and weathering-related magnetite-, hematite-, and goethite-rich iron ore: *Economic Geology*, v. 105, p. 917-945.

Angerer, T., Hagemann, S. G., and Danyushevsky, L. V., 2012, Geochemical evolution of the Banded Iron Formation-hosted high-grade iron ore system in the Koolyanobbing greenstone belt, Western Australia: *Economic Geology*, v. 107, p. 599-644.

Drake, J. R., 1972, The structure and petrology of banded iron formations at Mt. Windarra, Western Australia: Unpublished BSc (Hons) Thesis thesis, University of Western Australia, 62pp p.

Duuring, P., and Hagemann, S. G., 2010, High-grade iron mineralisation at the Beebyn deposit, Weld Range, Western Australia: 5th International Archean Symposium: Abstracts, Perth, Western Australia, 2010, p. 291-293.

---, 2011, Contrasting styles of high-grade iron mineralisation at Weld Range, Western Australia, *Iron Ore 2011: Perth, The Australasian Institute of Mining and Metallurgy*, p. 87-92.

---, 2012, Leaching of silica bands and concentration of magnetite in Archean BIF by hypogene fluids: Beebyn Fe ore deposit, Yilgarn Craton, Western Australia. *Mineralium Deposita*. doi: 10.1007/s00126-012-0428-1.

---, 2012, Genesis of superimposed hypogene and supergene Fe orebodies in BIF at the Madoonga deposit, Yilgarn Craton, Western Australia. *Mineralium Deposita*. doi: 10.1007/s00126-012-0429-0.

Duuring P, Hagemann SG, Novikova J, Cudahy T, Laukamp C., 2012, Targeting iron ore in Banded Iron Formation using ASTER data: Weld Range greenstone belt, Yilgarn Craton, Western Australia. *Economic Geology*, v. 107, p. 585-597.

Greentree, M. R., and Lord, D., 2007, Iron mineralization in the Yilgarn craton and future potential, in Bierlein, F. P., and Knox-Robinson, C. M., eds., *Proceedings of Geoconferences (WA) Inc Kalgoorlie '07 Conference: Kalgoorlie*, p. 70-73.

Guarin, C. P., Angerer, T., Maund, N. H., Cowan, D. R., and Hagemann, S., 2009, The K Deeps Magnetite Mineralisation at Koolyanobbing, Western Australia: *Iron Ore 2009 Proceedings*, p. 95-104.

Lascelles, D. F., 2006, The Genesis of the Hope Downs Iron Ore Deposit, Hamersley Province, Western Australia: *Economic Geology*, v. 101, p. 1359-1376.

Lascelles, D. F., 2006, The Mount Gibson banded iron formation-hosted magnetite deposit: Two distinct processes for the origin of high-grade iron ore: *Economic Geology*, v. 101, p. 651-666.

Lascelles, D. F., 2009, Microplaty hematite in the Yilgarn: The Wiluna West Iron Ore Deposit, in Williams, P. J., ed., *Proceedings of the 10th Biennial SGA Meeting of the Society for Geology Applied to Mineral Deposits: Townsville, Australia*.

Maskell, A., 2010, Multi-stage structural controls and hypogene alteration of high-grade Fe-mineralisation at Matthew Ridge prospect, Jack Hills greenstone belt, Yilgarn craton: Unpublished Honours thesis, The University of Western Australia, 50 p.

Maskell, A., Duuring, P., and Hagemann, S. G., submitted, Hydrothermal alteration events controlling magnetite-rich iron ore at the Matthew Ridge prospect, Jack Hills greenstone belt, Yilgarn Craton: *Australian Journal of Earth Sciences*.

Miles Keith, R., 1953, Wilgie Mia, Weld range, iron ore: *Proceedings Empire Mining and Metallurgical Congress*, v. 1, p. 242-244.

Pilbara Craton, Western Australia

To be populated.

Africa

Cope, I. L., Wilkinson, J. J., Boyce, A. J., Chapman, J. B., Herrington, R. J., and Harris, C. J., 2008, Genesis of the Pic de Fon Iron Oxide Deposit, Simandou Range, Republic of Guinea, West Afrika, in Hagemann, S., Rosière, C. A., Gutzmer, J., and Beukes, N. J., eds., *Banded Iron Formation-related High-grade Iron Ore. Reviews in Economic Geology*, v. 15, p. 339-360.

Evans, D. A. D., Beukes, N. J., Gutzmer, J., and Kirschvink, J. L., 2002, Supergene and hypogene hematization events associated with Paleoproterozoic iron and manganese ores, South Africa, *Geological Society of America, 2002 annual meeting.*, v. 34, p. 382.

Gutzmer, J., Beukes, N. J., de Kock, M. O., and Netshiozwi, S. T., 2005, Origin of High-Grade Iron Ores at the Thabazimbi Deposit, South Africa, in AusIMM, ed., Proceedings Iron Ore Conference 2005, 19 - 21 September 2005, Freemantle, Western Australia, p. 57-65.

Brazil

Barbour, A. P., 1973, Distribution of Phosphorus in the Iron Ore Deposits of Itabira, Minas Gerais, Brazil: *Economic Geology*, v. 68, p. 52-64.

Cabral, A. R., Lehmann, B., Galbiatti, H. F., and Filho, O. G. R., 2006, Evidence for metre-scale variations in hematite composition within the Palaeoproterozoic Itabira Iron Formation, Minas Gerais, Brazil: *Mineral Mag*, v. 70, p. 591-602.

Cabral, A. R., Rocha Filho, O. G., and Jones, R. D., 2003, Hydrothermal origin of soft hematite ore in the Quadrilátero Ferrífero of Minas Gerais, Brazil: petrographic evidence from the Gongo Soco iron ore deposit: *Applied Earth Science*, v. 112, p. 279-286.

Cabral, A. R., Rocha Filho, O. G., and Jones, R. D., 2008, Mercury in itabirite-hosted soft hematite ore in the Quadrilátero Ferrífero of Minas Gerais: *Journal of Geochemical Exploration*, v. 96, p. 69-76.

Dorr, J. V. N., 1964, Supergene iron ores of Minas Gerais, Brazil: *Economic Geology*, v. 59, p. 1203-1240.

---, 1965, Nature and origin of the high- grade hematite ores of Minas Gerais, Brazil: *Economic Geology*, v. 60, p. 1-46.

[Figueiredo e Silva, R. C., Lobato, L. M., and Rosiere, C. A., 2008, A Hydrothermal Origin for the Jaspilite-Hosted Giant Sierra Norte Deposits in the Cajajas Mineral Province, Para State, Brazil, in Hagemann, S., Rosière, C. A., Gutzmer, J., and Beukes, N. J., eds., *Banded Iron Formation-related High-grade Iron Ore*, v. 15, *Society of Economic Geologists*, p. 255-290.](#)

Galbiatti, H. F., Fonseca, M. A., Pereira, M. C., and Polônia, J. C., 2007, Structural control of Au–Pd mineralization (Jacutinga): An example from the Cauê Mine, Quadrilátero Ferrífero, Brazil: *Ore Geology Reviews*, v. 32, p. 614-628.

Hoefs, J., Müller, G., and Schuster, A. K., 1982, Polymetamorphic relations in iron ores from the Iron Quadrangle, Brazil: The correlation of oxygen isotope variations with deformation history: *Contributions to Mineralogy and Petrology*, v. 79, p. 241-251.

Klein, C., and Ladeira, E. A., 2000, Geochemistry and petrology of some Proterozoic banded iron-formations of the Quadrilátero Ferrífero, Minas Gerais, Brazil: *Economic Geology and the Bulletin of the Society of Economic Geologists*, v. 95, p. 405-427.

Lagoeiro, L. E., 1998, Transformation of magnetite to hematite and its influence on the dissolution of iron oxide minerals: *Journal of Metamorphic Geology*, v. 16, p. 415-423.

Luders, V., Romer, R. L., Cabral, A. R., Schmidt, C., Banks, D. A., and Schneider, J., 2005, Genesis of itabirite-hosted Au-Pd-Pt-bearing hematite-(quartz) veins, Quadrilátero Ferrífero, Minas Gerais, Brazil: constraints from fluid inclusion infrared microthermometry, bulk crush-leach analysis and U-Pb systematics: *Mineralium Deposita*, v. 40, p. 289-306.

Olivo, G. R., Gauthier, M., Williams-Jones, A. E., and Levesque, M., 2001, The Au-Pd mineralization at the Conceicao iron mine, Itabira district, southern Sao Francisco craton, Brazil: An example of a Jacutinga-type deposit: *Economic Geology and the Bulletin of the Society of Economic Geologists*, v. 96, p. 61-74.

Pereira, A., Fabris, J., Rios, F., Rosière, C., da Costa Couceiro, P., Ferreira, F., and de Menezes, L., 2010, Hematite from a mining area in the east border of Quadrilátero Ferrífero, Minas Gerais, Brazil: Hyperfine Interactions, v. 195, p. 69-76.

Pires, F. R. M., 2003, Distribution of hard hematite ore at the Quadrilátero Ferrífero, Minas Gerais, Brazil and its possible genetic significance: *Transactions of the Institution of Mining and Metallurgy Section B-Applied Earth Science*, v. 112, p. B31-B37.

Rosière, C. A., and Chemale Jr, F., 2000, Brazilian iron formations and their geological setting.

Rosiere, C. A., and Rios, F. J., 2004, The origin of hematite in high-grade iron ores based on infrared microscopy and fluid inclusion studies: The example of the conceicao mine, Quadrilátero Ferrífero, Brazil: *Economic Geology and the Bulletin of the Society of Economic Geologists*, v. 99, p. 611-624.

Rosiere, C. A., Siemes, H., Quade, H., Brokmeier, H. G., and Jansen, E. M., 2001, Microstructures, textures and deformation mechanisms in hematite: *Journal of Structural Geology*, v. 23, p. 1429-1440.

Rosiere, C. A., Spier, C. A., Rios, F. J., and Suckau, V. E., 2008, The Itabirites of the Quadrilátero Ferrífero and Related High-Grade Iron Ore Deposits: An Overview: *Banded Iron Formation-Related High-Grade Iron Ore*, v. 15, p. 223-254.

Selmi, M., Lagoeiro, L. E., and Endo, I., 2009, Geochemistry of hematite and itabirite, Quadrilátero Ferrífero, Brazil: *Rem: Revista Escola de Minas*, v. 62, p. 35-43.

Silva, R. C. F. E., Hagemann, S. G., Lobato, L. M., Rosiere, C. A., and Vennemann, T., 2008, Iron oxide paragenesis, quartz vein chronology and hydrothermal fluid evolution at the giant North Range Carajas iron deposits in Brazil, 2008.

Silva, R. C. F. E., Lobato, L. M., Rosiere, C. A., Hagemann, S., Zucchetti, M., Baars, F. J., Morais, R., and Andrade, I., 2008, A Hydrothermal Origin for the Jaspilite-Hosted, Giant Serra Norte Iron Ore Deposits in the Carajas Mineral Province, Para State, Brazil: *Banded Iron Formation-Related High-Grade Iron Ore*, v. 15, p. 255-290.

Spier, C. A., de Oliveira, S. M. B., and Rosiere, C. A., 2003, Geology and geochemistry of the Aguas Claras and Pico Iron Mines, Quadrilátero Ferrífero, Minas Gerais, Brazil: *Mineralium Deposita*, v. 38, p. 751-774.

Spier, C. A., de Oliveira, S. M. B., Rosiere, C. A., and Ardisson, J. D., 2008, Mineralogy and trace-element geochemistry of the high-grade iron ores of the Aguas Claras Mine and comparison with the Capao Xavier and Tamandua iron ore deposits, Quadrilhero Ferrifero, Brazil: *Mineralium Deposita*, v. 43, p. 229-254.

Varajao, C. A. C., Bruand, A., Ramandaidou, E. R., and Gilkes, R. J., 2002, Microporosity of BIF hosted massive hematite ore, Iron Quadrangle, Brazil: *Anais Da Academia Brasileira De Ciencias*, v. 74, p. 113-126.

Other topics, including glossary of terms

[Duuring P, Hagemann SG, Novikova J, Cudahy T, Laukamp C., 2012, Targeting iron ore in Banded Iron Formation using ASTER data: Weld Range greenstone belt, Yilgarn Craton, Western Australia. *Economic Geology*, v. 107, p. 585-597.](#)

Gutzmer, J., Chisonga, C. B., Beukes, N. J., and Mukhopadhyay, J., 2008, The Geochemistry of Iron Formation-Hosted High-Grade Hematite-Martite Iron Ores, in Hagemann, S. G., Rosière, C. A., Gutzmer, J., and Beukes, N. J., eds., *Banded Iron Formation-related High-grade Iron Ore. Reviews in Economic Geology*, v. 15, p. 157-183.

Gutzmer, J., Mukhopadhyay, J., Beukes, N. J., Pack, A., Hayashi, K., and Sharp, Z. D., 2006, Oxygen isotope composition of hematite and genesis of high-grade BIF-hosted iron ores: *Geological Society of America Memoir*, v. 198, p. 257-268.

[**Hagemann, S., Rosiere, C. A., Gutzmer, J., and Beukes, N. J., 2008, Glossary of terms Banded Iron Formation-related high-grade iron ore: *Banded Iron Formation-Related High-Grade Iron Ore*, v. 15, p. 411-414.**](#)

Lagoeiro, L. E., 1998, Transformation of magnetite to hematite and its influence on the dissolution of iron oxide minerals: *Journal of Metamorphic Geology*, v. 16, p. 415-423.

Morris, R. C., and Fletcher, A. B., 1987, Increased solubility of quartz following ferrous-ferric iron reactions: *Nature*, p. 250-252.

Otake, T., Wesolowski, D. J., Anovitz, L. M., Allard, L. F., and Ohmoto, H., 2007, Experimental evidence for non-redox transformations between magnetite and hematite under H₂-rich hydrothermal conditions: *Earth and Planetary Science Letters*, v. 257, p. 60-70.

Ramanaidou, E., Wells, M., Belton, D., Verrall, M., and Ryan, C., 2008, Mineralogical and Microchemical Methods for the Characterization of High-Grade Banded Iron Formation-Derived Iron Ore in Hagemann, S., Rosière, C. A., Gutzmer, J., and Beukes, N. J., eds., *Banded Iron Formation-related High-grade Iron Ore*, v. 15, Society of Economic Geologists, p. 129-156.

Rosiere, C. A., Siemes, H., Quade, H., Brokmeier, H. G., and Jansen, E. M., 2001, Microstructures, textures and deformation mechanisms in hematite: *Journal of Structural Geology*, v. 23, p. 1429-1440.